

A GRAY WHALE SIGHTING OFF HAWAI'I ISLAND: THE FIRST RECORD FOR THE CENTRAL TROPICAL PACIFIC

Robin W. Baird

Cascadia Research Collective, 218 ½ W. 4th Avenue, Olympia, Washington, 98501
rwbaird@cascadiaresearch.org

Jesse James

78-6629 Alii Drive, Kailua-Kona, Hawai'i 96740

Chad Mata

P.O. Box 162, Honaunau, Hawai'i, 96726

On February 2, 2022, a gray whale (*Eschrichtius robustus*) was sighted during a commercial whale watching trip in shallow (<100 m) water north of Keāhole Point, Hawai'i Island (approximately 19.73°N, 156.07°W). This is the first record of this species in Hawaiian waters and in the central tropical Pacific. Video footage was taken with a cell phone during the sighting. Based on size noted in the field and the amount of barnacle scarring visible, this individual is likely a juvenile (A. Perez, pers. comm.). An examination of the video footage suggests the body condition of the individual may be compromised (A.L. Bradford, pers. comm.), but the resolution of the video precluded a more precise determination (i.e., fair versus poor; Bradford et al. 2012). Preliminary comparisons to the Cascadia Research Collective gray whale catalog from the eastern North Pacific (Calambokidis et al. 2019) have been undertaken with no matches to date. Video stills will be sent to other regional gray whale catalogs for comparison.

Gray whales are primarily restricted to coastal waters during breeding, feeding, and migration, and this individual would have had to have crossed over 3,000 km of deep, open-ocean waters from the nearest area where this species is known to seasonally occupy, off the Aleutian Islands. The eastern North Pacific population breeds in shallow lagoons along the west coast of Baja California, Mexico, and feeds in shallow coastal waters from northern California north along the coasts of Oregon, Washington, British Columbia, Alaska, and Russia (Heide-Jorgensen et al. 2012; Lagerquist et al. 2019; Urbán et al. 2021). Migrating gray whales in the eastern North Pacific appear to largely remain on the continental shelf to and from their feeding grounds to the north (Lagerquist et al. 2019; Urbán et al. 2021). In the western North Pacific, gray whales feed in shallow waters off Sakhalin Island and the Kamchatka Peninsula in Russia, and historically would winter off southern China (Weller et al. 2008; Wang et al. 2015). More recent evidence shows that at least some whales from the western Pacific migrate east past the Aleutians and may cross through the deep waters of the Gulf of Alaska to the coast of British Columbia and Washington state before heading south along the continental shelf to breeding grounds in Mexico (Mate et al. 2015). Long-distance and open ocean movements of gray whales have been documented several times in recent years, including an individual photographed and biopsy sampled off Namibia (Elwen and Gridley 2013; Hoelzel et al. 2021), and a different individual photographed in the Mediterranean off both Israel and Spain (Scheinin et al. 2011).

Prior to this sighting seven species of baleen whales had been documented in Hawaiian waters, a few of which are rarely seen (Baird 2016). While humpback whales (*Megaptera novaeangliae*)

are abundant in nearshore Hawaiian waters in the winter, other baleen whale species primarily use pelagic waters (e.g., Bryde's whales, *Balaenoptera edeni*) and are only infrequently seen (e.g., fin whales, *B. physalus*, and sei whales, *B. borealis*). Blue whales (*Balaenoptera musculus*) have only been seen once in Hawaiian waters, and North Pacific right whales (*Eubalaena japonica*) and minke whales (*B. acutorostrata*) on just a few occasions each (Bradford et al. 2017; Herman et al. 1980; Salden and Mickelsen 1999), although blue and minke whales are regularly detected acoustically in Hawaiian waters (Norris et al. 2012; Oswald et al. 2011; Stafford et al. 2001; Thompson and Friedl 1982). The February 2022 sighting suggests that there may be value in assessing existing acoustic archives from hydrophones in Hawaiian waters for the presence of gray whale calls (e.g., Guazzo et al. 2017).

Boat-based research efforts on cetaceans in Hawaiian waters began in the late 1960s (Norris and Dohl 1980), and for the last 35 years multiple research groups have worked with cetaceans off the main Hawaiian Islands each year. Despite this extensive effort, this first sighting of a gray whale in Hawai'i was made during a commercial whale watching trip, another demonstration of the value of community science to understanding whales and dolphins in Hawaiian waters.

Acknowledgements

We thank Hawaii Oceanic for the platform from which this sighting was made, Jordan Lerma for video processing, Alie Perez for taking images of the whale through the Cascadia Research gray whale catalog, Amanda Bradford for an assessment of body condition, and Amanda Bradford, Annette Harnish, Michaela Kratofil, Sabre Mahaffy, and Alie Perez for reviews of the manuscript.

Literature Cited

- Baird, R.W. 2016. The lives of Hawai'i's dolphins and whales: natural history and conservation. University of Hawai'i Press.
- Bradford, A.L., D.W. Weller, A.E. Punt, Y.V. Ivashenko, A.M. Burdin, G.R. VanBlaricom, and R.L. Brownell Jr. 2012. Leaner leviathans: body condition variation in a critically endangered whale population. *Journal of Mammalogy* 93:251-266. 10.1644/11-MAMM-A-091.1.
- Bradford, A.L., K.A. Forney, E.M. Oleson, and J. Barlow. 2017. Abundance estimates of cetaceans from a line-transect survey within the U.S. Hawaiian Islands Exclusive Economic Zone. *Fishery Bulletin* 115:129-142. 10.7755/FB.115.2.1
- Calambokidis, J., A. Pérez, and J. Laake. 2019. Updated analysis of abundance and population structure of seasonal gray whales in the Pacific Northwest, 1996-2017. Final Report to NOAA, Seattle, Washington. pp. 1-72.
- Elwen, S.H., and T. Gridley. 2013. Gray whale (*Eschrichtius robustus*) sighting in Namibia (SE Atlantic) – first record for southern hemisphere. IWC SC/65a/BRG30. International Whaling Commission.

- Guazzo, R.A., T.A. Helble, G.L. D'Spain, D.W. Weller, S.M. Wiggins, and J.A. Hildebrand. 2017. Migratory behavior of eastern North Pacific gray whales tracked using a hydrophone array. *PLoS ONE* 12(10):e0185585.
- Heide-Jorgensen, M.P., K.L. Laidre, D. Litovka, M. Villum Jensen, J.M. Grebmeier, and B.I. Sirenko. 2012. Identifying gray whale (*Eschrichtius robustus*) foraging grounds along the Chukotka Peninsula, Russia, using satellite telemetry. *Polar Biology* 35:1035-1045.
- Herman, L.M., C.S. Baker, P.H. Forestell, and R.C. Antionja. 1980. Right whale *Balaena glacialis* sightings near Hawaii: a clue to the wintering grounds? *Marine Ecology Progress Series* 2:271-275.
- Hoelzel, A.R., F. Sarigol, T. Gridley, and S.H. Elwen. 2021. Natal origin of Namibian grey whale implies new distance record for in-water migration. *Biology Letters* 17: 20210136. 10.1098/rsbl.2021.0136.
- Lagerquist, B.A., D.M. Palacios, M.H. Winsor, L.M. Irvine, T.M. Follett, and B.R. Mate. 2019. Feeding home ranges of Pacific coast feeding group gray whales. *Journal of Wildlife Management* 83:925-937. 10.1002/jwmg.21642.
- Mate, B.R., V.Y. Ilyashenko, A.L. Bradford, V.V. Vertyankin, G.A. Tsidulko, V.V. Rozhnov, and L.M. Irvine. 2015. Critically endangered western gray whales migrate to the eastern North Pacific. *Biology Letters* 11:20150071. 10.1098/rsbl.2015.0071
- Norris, K.S., and T.P. Dohl. 1980. Behavior of the Hawaiian spinner dolphin, *Stenella longirostris*. *Fishery Bulletin* 77:821-849.
- Norris, T., S. Martin, L. Thomas, T. Yack, J.N. Oswald, E.M. Nosal, and V. Janik. 2012. Acoustic ecology and behavior of minke whales in the Hawaiian and Marianas Islands: localization, abundance estimation, and characterization of minke whale "boings". In *The effects of noise on aquatic life* (pp. 149-153). Springer, New York, NY.
- Oswald, J.N., W.W.L. Au, and F. Duennebieer. 2011. Minke whale (*Balaenoptera acutorostrata*) boings detected at the Station ALOHA cabled observatory. *Journal of the Acoustical Society of America* 129:3353-3360.
- Salden, D.R., and J. Mickelsen. 1999. Rare sighting of a North Pacific right whale (*Eubalaena glacialis*) in Hawai'i. *Pacific Science* 53:341-345.
- Scheinin, A.P., D. Kerem, C.D. Macleod, M. Gazo, C.A. Chicote, and M. Castellote. 2011. Gray whale (*Eschrichtius robustus*) in the Mediterranean Sea: anomalous event or early sign of climate-driven distribution change? *Marine Biodiversity Records* 4:e28. 10.1017/S1755267211000042.

- Stafford, K.M., S.L. Nieukirk, and C.G. Fox. 2001. Geographic and seasonal variation of blue whale calls in the North Pacific. *Journal of Cetacean Research and Management* 3:65-76.
- Thompson, P.O. and W.A. Friedl. 1982. A long term study of low frequency sounds from several species of whales off Oahu, Hawaii. *Cetology* 45:1-19.
- Urbán R, J., E. Jiménez-López, H.M. Guzmán, and L. Vilorio-Gómora. 2021. Migratory behavior of an eastern North Pacific gray whale from Baja California Sur to Chirikov Basin, Alaska. *Frontiers in Marine Science* 10.3380/fmars.2021.619290.
- Wang, X., M. Xu, F. Wu, D.W. Weller, X. Miao, A.R. Lang, and Q. Zhu. 2015. Insights from a gray whale (*Eschrichtius robustus*) bycaught in the Taiwan Strait off China in 2011. *Aquatic Mammals* 41:327-332.
- Weller, D.W., A.L. Bradford, H. Kato, T. Bando, S. Otani, A.M. Burdin, and R.L. Brownell Jr. 2008. A photographic match of a western gray whale between Sakhalin Island, Russia, and Honshu, Japan: the first link between the feeding ground and a migratory corridor. *Journal of Cetacean Research and Management* 10:89-91.



Figure 1. Images taken from video of a gray whale documented north of Keāhole Point, Hawai‘i, on February 2, 2022. The Blue Ocean Mariculture kanpachi (*Seriola rivoliana*) farm is visible in the background of the lower two images.